

CAMPBELL BIOLOGY DNA REPLICATION KEY TERMS

CAMPBELL BIOLOGY DNA REPLICATION KEY TERMS: A COMPREHENSIVE GUIDE

CAMPBELL BIOLOGY DNA REPLICATION KEY TERMS ARE FUNDAMENTAL TO UNDERSTANDING ONE OF LIFE'S MOST CRUCIAL PROCESSES: THE DUPLICATION OF GENETIC MATERIAL. THIS ARTICLE DELVES DEEP INTO THE CORE CONCEPTS AND TERMINOLOGY PRESENTED IN CAMPBELL BIOLOGY'S TREATMENT OF DNA REPLICATION, PROVIDING A DETAILED EXPLORATION OF THE ENZYMES, MECHANISMS, AND REGULATORY ASPECTS INVOLVED. WE WILL UNRAVEL THE INTRICATE DANCE OF DNA POLYMERASE, HELICASE, LIGASE, AND OTHER ESSENTIAL PLAYERS, ILLUMINATING HOW THE CELL METICULOUSLY COPIES ITS GENOME WITH REMARKABLE FIDELITY. UNDERSTANDING THESE KEY TERMS IS PARAMOUNT FOR STUDENTS OF BIOLOGY, RESEARCHERS, AND ANYONE SEEKING A PROFOUND GRASP OF MOLECULAR GENETICS.

TABLE OF CONTENTS

THE DOUBLE HELIX: A FOUNDATION FOR REPLICATION

INITIATION OF DNA REPLICATION: GETTING STARTED

ELONGATION: BUILDING NEW DNA STRANDS

TERMINATION OF DNA REPLICATION: COMPLETING THE PROCESS

ENZYMES AND PROTEINS: THE MOLECULAR MACHINERY

ACCURACY AND REPAIR: ENSURING FIDELITY

PROKARYOTIC VS. EUKARYOTIC DNA REPLICATION: KEY DIFFERENCES

TELOMERES AND TELOMERASE: THE ENDS OF THE CHROMOSOMES

THE DOUBLE HELIX: A FOUNDATION FOR REPLICATION

BEFORE DIVING INTO THE MECHANICS OF DNA REPLICATION, IT'S ESSENTIAL TO REVISIT THE ICONIC DOUBLE HELIX STRUCTURE. AS ELUCIDATED BY WATSON AND CRICK, DNA IS A POLYMER OF NUCLEOTIDES, EACH CONSISTING OF A DEOXYRIBOSE SUGAR, A PHOSPHATE GROUP, AND ONE OF FOUR NITROGENOUS BASES: ADENINE (A), GUANINE (G), CYTOSINE (C), AND THYMINE (T). THESE NUCLEOTIDES ARE LINKED BY PHOSPHODIESTER BONDS, FORMING A SUGAR-PHOSPHATE BACKBONE. THE TWO STRANDS OF THE HELIX RUN ANTIPARALLEL, MEANING THEIR 5' AND 3' ENDS ARE ORIENTED IN OPPOSITE DIRECTIONS.

THE KEY TO DNA REPLICATION LIES IN THE COMPLEMENTARY BASE PAIRING RULES: ADENINE ALWAYS PAIRS WITH THYMINE (A-T) VIA TWO HYDROGEN BONDS, AND GUANINE ALWAYS PAIRS WITH CYTOSINE (G-C) VIA THREE HYDROGEN BONDS. THIS INHERENT COMPLEMENTARITY ENSURES THAT EACH STRAND SERVES AS A PRECISE TEMPLATE FOR THE SYNTHESIS OF A NEW, IDENTICAL STRAND DURING REPLICATION. THIS SEMI-CONSERVATIVE MECHANISM, WHERE EACH NEW DNA MOLECULE CONSISTS OF ONE ORIGINAL STRAND AND ONE NEWLY SYNTHESIZED STRAND, IS A CORNERSTONE OF MODERN GENETICS.

INITIATION OF DNA REPLICATION: GETTING STARTED

DNA REPLICATION DOES NOT BEGIN RANDOMLY; IT IS A HIGHLY REGULATED PROCESS INITIATED AT SPECIFIC SITES WITHIN THE GENOME. THESE DESIGNATED STARTING POINTS ARE KNOWN AS ORIGINS OF REPLICATION. IN PROKARYOTES, SUCH AS BACTERIA, THERE IS TYPICALLY A SINGLE ORIGIN OF REPLICATION ON THEIR CIRCULAR CHROMOSOME, WHILE EUKARYOTES, WITH THEIR LARGER AND MORE COMPLEX LINEAR CHROMOSOMES, POSSESS MULTIPLE ORIGINS OF REPLICATION ALONG EACH CHROMOSOME. THIS MULTI-ORIGIN STRATEGY IS CRUCIAL FOR EFFICIENTLY REPLICATING THE VAST AMOUNT OF GENETIC MATERIAL IN EUKARYOTIC CELLS WITHIN A REASONABLE TIMEFRAME.

THE PROCESS OF INITIATION INVOLVES THE BINDING OF INITIATOR PROTEINS TO THE ORIGIN OF REPLICATION. THESE PROTEINS RECRUIT OTHER FACTORS THAT HELP TO UNWIND THE DNA DOUBLE HELIX, SEPARATING THE TWO PARENTAL STRANDS. THIS UNWINDING CREATES A REPLICATION BUBBLE, WITH TWO REPLICATION FORKS EXTENDING OUTWARDS IN OPPOSITE DIRECTIONS. EACH REPLICATION FORK IS A Y-SHAPED STRUCTURE WHERE THE PARENTAL DNA IS ACTIVELY BEING UNWOUND AND NEW DNA STRANDS ARE BEING SYNTHESIZED.

ELONGATION: BUILDING NEW DNA STRANDS

ONCE THE REPLICATION FORKS ARE ESTABLISHED, THE PROCESS OF ELONGATION BEGINS. THIS IS WHERE THE BULK OF DNA SYNTHESIS OCCURS, WITH ENZYMES METICULOUSLY ADDING NUCLEOTIDES TO THE GROWING NEW STRANDS. THE PRIMARY ENZYME RESPONSIBLE FOR THIS TASK IS DNA POLYMERASE. HOWEVER, DNA POLYMERASE CANNOT INITIATE DNA SYNTHESIS ON A BARE TEMPLATE STRAND; IT REQUIRES A PRE-EXISTING SHORT STRAND OF NUCLEOTIDES, KNOWN AS A PRIMER, TO ATTACH NEW NUCLEOTIDES TO. PRIMERS ARE SYNTHESIZED BY AN ENZYME CALLED PRIMASE.

DNA POLYMERASE ADDS NEW NUCLEOTIDES TO THE 3' END OF THE GROWING STRAND. THIS DIRECTIONALITY IS CRITICAL AND LEADS TO A KEY CHARACTERISTIC OF DNA REPLICATION: THE SYNTHESIS OF THE TWO NEW STRANDS OCCURS DIFFERENTLY. THE STRAND SYNTHESIZED IN THE 5' TO 3' DIRECTION, CONTINUOUSLY MOVING TOWARDS THE REPLICATION FORK, IS CALLED THE LEADING STRAND. IN CONTRAST, THE OTHER STRAND, SYNTHESIZED IN THE OPPOSITE DIRECTION (AWAY FROM THE REPLICATION FORK), IS FORMED DISCONTINUOUSLY IN SHORT FRAGMENTS CALLED OKAZAKI FRAGMENTS. THIS STRAND IS KNOWN AS THE LAGGING STRAND.

THE LEADING STRAND SYNTHESIS

THE LEADING STRAND IS SYNTHESIZED CONTINUOUSLY BECAUSE DNA POLYMERASE CAN MOVE IN THE 5' TO 3' DIRECTION, FOLLOWING THE UNWINDING OF THE DNA HELIX AT THE REPLICATION FORK. ONLY A SINGLE PRIMER IS NEEDED AT THE ORIGIN OF REPLICATION TO INITIATE LEADING STRAND SYNTHESIS. AS THE HELIX UNWINDS, DNA POLYMERASE ADDS NUCLEOTIDES ONE BY ONE, CREATING A SEAMLESS NEW STRAND THAT IS COMPLEMENTARY TO THE TEMPLATE STRAND.

THE LAGGING STRAND SYNTHESIS

THE LAGGING STRAND PRESENTS A MORE COMPLEX CHALLENGE. BECAUSE DNA POLYMERASE CAN ONLY SYNTHESIZE IN THE 5' TO 3' DIRECTION, AND THE LAGGING STRAND TEMPLATE IS ORIENTED IN THE 3' TO 5' DIRECTION RELATIVE TO THE REPLICATION FORK'S MOVEMENT, SYNTHESIS MUST OCCUR IN SHORT BURSTS. PRIMASE LAYS DOWN MULTIPLE RNA PRIMERS ALONG THE LAGGING STRAND TEMPLATE AS THE HELIX UNWINDS. DNA POLYMERASE THEN SYNTHESIZES SHORT DNA FRAGMENTS, CALLED OKAZAKI FRAGMENTS, STARTING FROM EACH PRIMER AND EXTENDING UNTIL IT REACHES THE PREVIOUS OKAZAKI FRAGMENT OR RUNS OUT OF TEMPLATE.

JOINING OKAZAKI FRAGMENTS

THE PRESENCE OF OKAZAKI FRAGMENTS ON THE LAGGING STRAND NECESSITATES A SUBSEQUENT STEP TO CREATE A CONTINUOUS DNA MOLECULE. AFTER DNA POLYMERASE SYNTHESIZES THE DNA PORTION OF AN OKAZAKI FRAGMENT, THE RNA PRIMER MUST BE REMOVED AND REPLACED WITH DNA. ENZYMES LIKE RNASE H ARE RESPONSIBLE FOR DEGRADING THE RNA PRIMER. THEN, ANOTHER TYPE OF DNA POLYMERASE FILLS THE GAP WITH DNA NUCLEOTIDES. FINALLY, THE ENZYME DNA LIGASE SEALS THE NICKS BETWEEN THE OKAZAKI FRAGMENTS AND THE NEWLY SYNTHESIZED DNA, FORMING A CONTINUOUS PHOSPHODIESTER BACKBONE.

TERMINATION OF DNA REPLICATION: COMPLETING THE PROCESS

TERMINATION OF DNA REPLICATION VARIES BETWEEN PROKARYOTES AND EUKARYOTES. IN PROKARYOTES, WITH THEIR CIRCULAR CHROMOSOMES, REPLICATION FORKS MEET AT A SPECIFIC TERMINATION REGION ON THE OPPOSITE SIDE OF THE ORIGIN. SPECIFIC PROTEINS BIND TO THIS REGION AND HALT THE PROGRESS OF THE REPLICATION FORKS, PREVENTING OVER-REPLICATION. THE TWO NEW CIRCULAR DNA MOLECULES ARE THEN SEPARATED.

IN EUKARYOTES, THE TERMINATION PROCESS IS MORE COMPLEX DUE TO MULTIPLE ORIGINS AND LINEAR CHROMOSOMES. WHEN REPLICATION FORKS MOVING FROM ADJACENT ORIGINS MEET, THE REMAINING DNA SEGMENTS ARE PROCESSED AND LIGATED. HOWEVER, A UNIQUE CHALLENGE ARISES AT THE VERY ENDS OF LINEAR CHROMOSOMES, KNOWN AS TELOMERES. THE LAGGING STRAND SYNTHESIS PROBLEM AT THE VERY END OF THE CHROMOSOME LEADS TO A PROGRESSIVE SHORTENING OF DNA WITH EACH ROUND OF REPLICATION. THIS ISSUE IS ADDRESSED BY A SPECIALIZED ENZYME.

ENZYMES AND PROTEINS: THE MOLECULAR MACHINERY

THE REMARKABLE FIDELITY AND EFFICIENCY OF DNA REPLICATION ARE ORCHESTRATED BY A COMPLEX MOLECULAR MACHINERY COMPRISING NUMEROUS ENZYMES AND PROTEINS. EACH COMPONENT PLAYS A DISTINCT AND VITAL ROLE IN UNWINDING THE DNA, SYNTHESIZING NEW STRANDS, AND ENSURING ACCURACY.

- **HELICASE:** THIS ENZYME UNWINDS THE DNA DOUBLE HELIX AT THE REPLICATION FORK BY BREAKING THE HYDROGEN BONDS BETWEEN COMPLEMENTARY BASE PAIRS.
- **SINGLE-STRAND BINDING PROTEINS (SSBs):** THESE PROTEINS BIND TO THE SEPARATED SINGLE STRANDS OF DNA, PREVENTING THEM FROM RE-ANNEALING AND PROTECTING THEM FROM DEGRADATION.
- **TOPOISOMERASE:** AS HELICASE UNWINDS THE DNA, IT CAN CAUSE SUPERCOILING AHEAD OF THE REPLICATION FORK. TOPOISOMERASES RELIEVE THIS STRAIN BY CUTTING AND REJOINING THE DNA STRANDS.
- **PRIMASE:** THIS ENZYME SYNTHESIZES SHORT RNA PRIMERS, PROVIDING A NECESSARY STARTING POINT FOR DNA POLYMERASE.
- **DNA POLYMERASE:** THE CENTRAL ENZYME OF REPLICATION, DNA POLYMERASE SYNTHESIZES NEW DNA STRANDS BY ADDING COMPLEMENTARY NUCLEOTIDES TO THE 3' END OF A GROWING STRAND OR PRIMER. THERE ARE DIFFERENT TYPES OF DNA POLYMERASES WITH SPECIALIZED FUNCTIONS, INCLUDING PROOFREADING AND REPAIR.
- **DNA LIGASE:** THIS ENZYME SEALS THE NICKS IN THE SUGAR-PHOSPHATE BACKBONE, JOINING OKAZAKI FRAGMENTS ON THE LAGGING STRAND AND COMPLETING THE SYNTHESIS OF CONTINUOUS DNA MOLECULES.
- **TELOMERASE:** IN EUKARYOTES, THIS ENZYME ADDS REPETITIVE DNA SEQUENCES TO THE ENDS OF LINEAR CHROMOSOMES (TELOMERES), COUNTERACTING THE SHORTENING THAT OCCURS DUE TO THE LAGGING STRAND SYNTHESIS PROBLEM.

ACCURACY AND REPAIR: ENSURING FIDELITY

DNA REPLICATION IS AN ASTONISHINGLY ACCURATE PROCESS, WITH ERROR RATES TYPICALLY IN THE RANGE OF ONE MISTAKE PER BILLION NUCLEOTIDES. THIS HIGH FIDELITY IS ACHIEVED THROUGH A COMBINATION OF MECHANISMS, PRIMARILY INVOLVING THE INTRINSIC PROOFREADING CAPABILITIES OF DNA POLYMERASE AND DEDICATED DNA REPAIR SYSTEMS. DNA POLYMERASE ITSELF HAS A 3' TO 5' EXONUCLEASE ACTIVITY, ALLOWING IT TO BACKTRACK, REMOVE INCORRECTLY INCORPORATED NUCLEOTIDES, AND REPLACE THEM WITH THE CORRECT ONES.

BEYOND THE POLYMERASE'S BUILT-IN ERROR CORRECTION, CELLS POSSESS SOPHISTICATED DNA REPAIR PATHWAYS THAT DETECT AND CORRECT ERRORS MISSED DURING REPLICATION. THESE PATHWAYS INCLUDE MISMATCH REPAIR, BASE EXCISION REPAIR, AND NUCLEOTIDE EXCISION REPAIR. THESE SYSTEMS WORK DILIGENTLY TO MAINTAIN THE INTEGRITY OF THE GENOME, PREVENTING THE ACCUMULATION OF MUTATIONS THAT COULD HAVE DETRIMENTAL EFFECTS ON CELLULAR FUNCTION AND ORGANISMAL HEALTH.

PROKARYOTIC VS. EUKARYOTIC DNA REPLICATION: KEY DIFFERENCES

WHILE THE FUNDAMENTAL PRINCIPLES OF DNA REPLICATION ARE CONSERVED ACROSS ALL LIFE FORMS, THERE ARE NOTABLE DIFFERENCES BETWEEN PROKARYOTIC AND EUKARYOTIC REPLICATION, REFLECTING THE DIFFERENCES IN GENOME STRUCTURE AND COMPLEXITY.

- **ORIGIN OF REPLICATION:** PROKARYOTES HAVE A SINGLE ORIGIN OF REPLICATION, WHEREAS EUKARYOTES HAVE MULTIPLE ORIGINS ON EACH LINEAR CHROMOSOME.
- **GENOME STRUCTURE:** PROKARYOTES TYPICALLY HAVE A CIRCULAR CHROMOSOME, WHILE EUKARYOTES HAVE LINEAR CHROMOSOMES.
- **REPLICATION RATE:** EUKARYOTIC REPLICATION IS GENERALLY SLOWER THAN PROKARYOTIC REPLICATION DUE TO THE GREATER COMPLEXITY AND SIZE OF THE EUKARYOTIC GENOME.
- **TELOMERES:** EUKARYOTES HAVE TELOMERES AT THE ENDS OF THEIR LINEAR CHROMOSOMES, WHICH REQUIRE THE ACTION OF TELOMERASE TO MAINTAIN THEIR LENGTH. PROKARYOTES, WITH THEIR CIRCULAR CHROMOSOMES, DO NOT HAVE THIS ISSUE.
- **DNA POLYMERASES:** WHILE BOTH HAVE MULTIPLE DNA POLYMERASES, THE SPECIFIC TYPES AND THEIR ROLES CAN DIFFER. EUKARYOTES HAVE A MORE EXTENSIVE SET OF DNA POLYMERASES INVOLVED IN VARIOUS ASPECTS OF DNA METABOLISM.

TELOMERES AND TELOMERASE: THE ENDS OF THE CHROMOSOMES

THE LINEAR NATURE OF EUKARYOTIC CHROMOSOMES PRESENTS A UNIQUE CHALLENGE FOR DNA REPLICATION. AS MENTIONED, THE LAGGING STRAND SYNTHESIS MECHANISM, WHICH INVOLVES DISCONTINUOUS OKAZAKI FRAGMENT SYNTHESIS, LEADS TO A PROGRESSIVE SHORTENING OF THE CHROMOSOME ENDS WITH EACH REPLICATION CYCLE. IF UNCHECKED, THIS SHORTENING WOULD EVENTUALLY LEAD TO THE LOSS OF ESSENTIAL GENETIC INFORMATION.

TELOMERES ARE SPECIALIZED REPETITIVE DNA SEQUENCES LOCATED AT THE ENDS OF EUKARYOTIC CHROMOSOMES. THESE SEQUENCES ACT AS PROTECTIVE CAPS, PREVENTING THE LOSS OF CODING DNA. THE ENZYME TELOMERASE PLAYS A CRUCIAL ROLE IN MAINTAINING TELOMERE LENGTH. TELOMERASE IS A REVERSE TRANSCRIPTASE THAT CARRIES ITS OWN RNA TEMPLATE AND CAN EXTEND THE 3' END OF THE PARENTAL DNA STRAND, PROVIDING A SUFFICIENT TEMPLATE FOR PRIMASE AND DNA POLYMERASE TO COMPLETE THE LAGGING STRAND SYNTHESIS AND COUNTERACT SHORTENING. THIS MAINTENANCE OF TELOMERE LENGTH IS VITAL FOR CELLULAR LONGEVITY AND PREVENTING GENOMIC INSTABILITY.

THE INTRICATE PROCESS OF DNA REPLICATION, AS DETAILED THROUGH THESE **CAMPBELL BIOLOGY DNA REPLICATION KEY TERMS**, UNDERSCORES THE ELEGANCE AND PRECISION OF MOLECULAR BIOLOGY. FROM THE INITIAL UNWINDING OF THE DOUBLE HELIX TO THE METICULOUS JOINING OF NEWLY SYNTHESIZED STRANDS AND THE PROTECTIVE MECHANISMS AT CHROMOSOME ENDS, EACH STEP IS A TESTAMENT TO THE FINELY TUNED MACHINERY OF LIFE. A SOLID UNDERSTANDING OF THESE CONCEPTS PROVIDES A CRITICAL FOUNDATION FOR FURTHER EXPLORATION IN GENETICS, MOLECULAR BIOLOGY, AND RELATED FIELDS.

FAQ

Q: WHAT IS THE PRIMARY FUNCTION OF HELICASE IN DNA REPLICATION ACCORDING TO

CAMPBELL BIOLOGY?

A: ACCORDING TO CAMPBELL BIOLOGY, THE PRIMARY FUNCTION OF HELICASE IN DNA REPLICATION IS TO UNWIND THE DNA DOUBLE HELIX AT THE REPLICATION FORK BY BREAKING THE HYDROGEN BONDS BETWEEN COMPLEMENTARY BASE PAIRS, THEREBY SEPARATING THE TWO PARENTAL STRANDS.

Q: EXPLAIN THE ROLE OF OKAZAKI FRAGMENTS IN DNA REPLICATION AS DESCRIBED IN CAMPBELL BIOLOGY.

A: CAMPBELL BIOLOGY EXPLAINS THAT OKAZAKI FRAGMENTS ARE SHORT, NEWLY SYNTHESIZED DNA FRAGMENTS THAT ARE FORMED ON THE LAGGING STRAND DURING DNA REPLICATION. BECAUSE DNA POLYMERASE CAN ONLY SYNTHESIZE DNA IN THE 5' TO 3' DIRECTION, THE LAGGING STRAND MUST BE SYNTHESIZED DISCONTINUOUSLY IN FRAGMENTS THAT ARE LATER JOINED TOGETHER.

Q: WHAT DOES CAMPBELL BIOLOGY MEAN BY "SEMI-CONSERVATIVE REPLICATION"?

A: CAMPBELL BIOLOGY DEFINES SEMI-CONSERVATIVE REPLICATION AS THE PROCESS WHERE EACH NEW DNA MOLECULE CONSISTS OF ONE ORIGINAL PARENTAL STRAND AND ONE NEWLY SYNTHESIZED DAUGHTER STRAND. THIS MEANS THE PARENTAL DOUBLE HELIX UNWINDS, AND EACH STRAND SERVES AS A TEMPLATE FOR THE SYNTHESIS OF A NEW COMPLEMENTARY STRAND.

Q: HOW DOES PROOFREADING BY DNA POLYMERASE CONTRIBUTE TO THE ACCURACY OF DNA REPLICATION, AS PER CAMPBELL BIOLOGY?

A: CAMPBELL BIOLOGY HIGHLIGHTS THAT DNA POLYMERASE POSSESSES A 3' TO 5' EXONUCLEASE ACTIVITY, WHICH ALLOWS IT TO PROOFREAD ITS WORK. IF AN INCORRECT NUCLEOTIDE IS ADDED TO THE GROWING DNA STRAND, THE POLYMERASE CAN BACKTRACK, REMOVE THE MISMATCHED NUCLEOTIDE, AND INSERT THE CORRECT ONE, SIGNIFICANTLY REDUCING THE ERROR RATE.

Q: WHAT IS THE SIGNIFICANCE OF TELOMERES AND TELOMERASE IN EUKARYOTIC DNA REPLICATION, ACCORDING TO CAMPBELL BIOLOGY?

A: CAMPBELL BIOLOGY EXPLAINS THAT TELOMERES ARE REPETITIVE DNA SEQUENCES AT THE ENDS OF EUKARYOTIC LINEAR CHROMOSOMES THAT PROTECT CODING REGIONS FROM LOSS DURING REPLICATION. TELOMERASE IS AN ENZYME THAT EXTENDS THESE TELOMERES, COUNTERACTING THE PROGRESSIVE SHORTENING THAT OCCURS WITH EACH ROUND OF REPLICATION DUE TO THE LAGGING STRAND SYNTHESIS PROBLEM.

Q: DIFFERENTIATE BETWEEN THE LEADING STRAND AND THE LAGGING STRAND SYNTHESIS DURING DNA REPLICATION, BASED ON CAMPBELL BIOLOGY'S EXPLANATION.

A: CAMPBELL BIOLOGY DIFFERENTIATES BY STATING THAT THE LEADING STRAND IS SYNTHESIZED CONTINUOUSLY IN THE 5' TO 3' DIRECTION, MOVING TOWARDS THE REPLICATION FORK. THE LAGGING STRAND, HOWEVER, IS SYNTHESIZED DISCONTINUOUSLY IN THE OPPOSITE DIRECTION (AWAY FROM THE REPLICATION FORK) IN SHORT OKAZAKI FRAGMENTS, WHICH ARE LATER JOINED BY DNA LIGASE.

Q: WHAT IS THE FUNCTION OF PRIMASE IN THE CONTEXT OF CAMPBELL BIOLOGY'S DISCUSSION ON DNA REPLICATION?

A: IN CAMPBELL BIOLOGY'S EXPLANATION, PRIMASE IS AN ENZYME THAT SYNTHESIZES SHORT RNA PRIMERS. THESE RNA PRIMERS ARE ESSENTIAL BECAUSE DNA POLYMERASE CANNOT INITIATE DNA SYNTHESIS DE NOVO; IT REQUIRES A PRE-EXISTING STRAND (THE PRIMER) TO ADD NUCLEOTIDES TO.

Campbell Biology Dna Replication Key Terms

Campbell Biology Dna Replication Key Terms

Related Articles

- [campbell biology climate change chapter exam us](#)
- [campbell biology chapter study guide us](#)
- [campbell biology developmental biology funding us](#)

[Back to Home](#)