

CLIMATE CHANGE EFFECTS MARINE LIFE

CLIMATE CHANGE EFFECTS MARINE LIFE ARE PROFOUND AND FAR-REACHING, IMPACTING ECOSYSTEMS FROM THE SURFACE TO THE DEEPEST TRENCHES. AS GLOBAL TEMPERATURES RISE AND ATMOSPHERIC CARBON DIOXIDE CONCENTRATIONS INCREASE, THE OCEANS ABSORB A SIGNIFICANT PORTION OF THIS EXCESS HEAT AND CO₂, LEADING TO A CASCADE OF DETRIMENTAL CHANGES. THESE ALTERATIONS ARE NOT ISOLATED INCIDENTS BUT INTERCONNECTED PHENOMENA THAT THREATEN THE BIODIVERSITY, ABUNDANCE, AND OVERALL HEALTH OF MARINE ORGANISMS AND THE VITAL SERVICES THEY PROVIDE. THIS ARTICLE DELVES INTO THE MULTIFACETED IMPACTS OF CLIMATE CHANGE ON MARINE ECOSYSTEMS, EXPLORING RISING OCEAN TEMPERATURES, OCEAN ACIDIFICATION, SEA-LEVEL RISE, ALTERED OCEAN CURRENTS, AND THE RESULTING CONSEQUENCES FOR MARINE SPECIES AND HABITATS.

TABLE OF CONTENTS

RISING OCEAN TEMPERATURES AND THEIR CONSEQUENCES
OCEAN ACIDIFICATION: THE SILENT THREAT
SEA-LEVEL RISE AND COASTAL MARINE HABITATS
CHANGES IN OCEAN CURRENTS AND NUTRIENT DISTRIBUTION
IMPACTS ON MARINE BIODIVERSITY AND FOOD WEBS
SPECIFIC VULNERABILITIES OF MARINE ECOSYSTEMS
CONSEQUENCES FOR HUMAN SOCIETIES

RISING OCEAN TEMPERATURES AND THEIR CONSEQUENCES

THE ABSORPTION OF EXCESS HEAT FROM THE ATMOSPHERE IS CAUSING A SIGNIFICANT AND ACCELERATING RISE IN OCEAN TEMPERATURES. THIS WARMING TREND HAS A DIRECT AND IMMEDIATE IMPACT ON MARINE ORGANISMS, MANY OF WHICH HAVE NARROW THERMAL TOLERANCE RANGES. AS WATERS BECOME WARMER, PHYSIOLOGICAL PROCESSES SUCH AS METABOLISM, GROWTH, AND REPRODUCTION CAN BE DISRUPTED, LEADING TO INCREASED STRESS AND MORTALITY.

CORAL REEFS ARE PARTICULARLY VULNERABLE TO RISING OCEAN TEMPERATURES. WHEN WATER TEMPERATURES EXCEED A CERTAIN THRESHOLD, CORALS EXPEL THE SYMBIOTIC ALGAE (ZOOXANTHELLAE) LIVING WITHIN THEIR TISSUES, A PROCESS KNOWN AS CORAL BLEACHING. WHILE BLEACHED CORALS ARE NOT DEAD, THEY ARE WEAKENED AND MORE SUSCEPTIBLE TO DISEASE. IF TEMPERATURES REMAIN ELEVATED FOR EXTENDED PERIODS, CORAL MORTALITY CAN BE WIDESPREAD, LEADING TO THE LOSS OF COMPLEX REEF STRUCTURES THAT SUPPORT A VAST ARRAY OF MARINE LIFE.

BEYOND CORAL REEFS, WARMING WATERS AFFECT A WIDE RANGE OF MARINE SPECIES. FISH POPULATIONS, FOR EXAMPLE, MAY MIGRATE TOWARDS COOLER POLAR REGIONS IN SEARCH OF SUITABLE HABITATS. THIS DISPLACEMENT CAN DISRUPT ESTABLISHED FOOD WEBS, IMPACT COMMERCIAL FISHERIES, AND ALTER THE DISTRIBUTION OF SPECIES, POTENTIALLY LEADING TO THE INTRODUCTION OF INVASIVE SPECIES INTO NEW AREAS. FURTHERMORE, INCREASED OCEAN TEMPERATURES CAN EXACERBATE EXISTING STRESSORS, SUCH AS POLLUTION AND OVERFISHING, MAKING MARINE ECOSYSTEMS LESS RESILIENT.

OCEAN ACIDIFICATION: THE SILENT THREAT

THE OCEAN ACTS AS A MASSIVE CARBON SINK, ABSORBING APPROXIMATELY 30% OF THE CARBON DIOXIDE (CO₂) RELEASED INTO THE ATMOSPHERE BY HUMAN ACTIVITIES. WHEN CO₂ DISSOLVES IN SEAWATER, IT FORMS CARBONIC ACID, WHICH THEN DISSOCIATES, RELEASING HYDROGEN IONS. THIS PROCESS LEADS TO A DECREASE IN THE PH OF SEAWATER, A PHENOMENON KNOWN AS OCEAN ACIDIFICATION. WHILE THE TERM "ACIDIFICATION" MIGHT SOUND ALARMING, IT REFERS TO A DECREASE IN ALKALINITY, MEANING THE OCEAN IS BECOMING LESS BASIC, NOT OUTRIGHT ACIDIC IN THE COMMON SENSE OF THE WORD.

THE MOST SIGNIFICANT IMPACTS OF OCEAN ACIDIFICATION ARE ON MARINE ORGANISMS THAT BUILD SHELLS AND SKELETONS MADE OF CALCIUM CARBONATE. THIS INCLUDES A DIVERSE GROUP OF ORGANISMS SUCH AS CORALS, SHELLFISH (OYSTERS, CLAMS, MUSSELS), PTEROPODS (SEA SNAILS), AND PHYTOPLANKTON. AS THE OCEAN BECOMES MORE ACIDIC, THE AVAILABILITY OF CARBONATE IONS, THE BUILDING BLOCKS FOR THESE STRUCTURES, DECREASES. THIS MAKES IT HARDER FOR THESE ORGANISMS TO GROW AND MAINTAIN THEIR SHELLS, AND IN SOME CASES, CAN EVEN CAUSE EXISTING SHELLS TO DISSOLVE.

THE CONSEQUENCES OF WIDESPREAD SHELL DISSOLUTION AND REDUCED CALCIFICATION ARE SEVERE. FOR CALCIFYING PLANKTON, WHICH FORM THE BASE OF MANY MARINE FOOD WEBS, A DECLINE IN THEIR POPULATIONS CAN HAVE CASCADING EFFECTS THROUGHOUT THE ECOSYSTEM. FOR COMMERCIALLY IMPORTANT SHELLFISH, ACIDIFICATION THREATENS AQUACULTURE AND WILD FISHERIES, IMPACTING ECONOMIES AND FOOD SECURITY. THE INTRICATE BALANCE OF MARINE ECOSYSTEMS, RELIANT ON THE HEALTH OF THESE CALCIFYING ORGANISMS, IS THUS UNDER SERIOUS THREAT FROM THIS INSIDIOUS CHANGE.

SEA-LEVEL RISE AND COASTAL MARINE HABITATS

GLOBAL WARMING IS CAUSING SEA LEVELS TO RISE THROUGH TWO PRIMARY MECHANISMS: THE THERMAL EXPANSION OF SEAWATER AS IT WARMS AND THE MELTING OF GLACIERS AND ICE SHEETS. THIS GRADUAL INCREASE IN SEA LEVEL POSES A SIGNIFICANT THREAT TO COASTAL MARINE HABITATS, WHICH ARE OFTEN HIGHLY PRODUCTIVE AND BIOLOGICALLY DIVERSE.

MANGROVE FORESTS AND SALT MARSHES, VITAL NURSERY GROUNDS FOR MANY FISH AND INVERTEBRATE SPECIES AND NATURAL BUFFERS AGAINST STORM SURGES, ARE PARTICULARLY VULNERABLE. AS SEA LEVELS RISE, THESE ECOSYSTEMS CAN BECOME INUNDATED, LEADING TO HABITAT LOSS IF THEY CANNOT MIGRATE INLAND. SEDIMENTATION PATTERNS CAN ALSO BE ALTERED, FURTHER IMPACTING THEIR HEALTH AND FUNCTIONALITY. COASTAL EROSION, EXACERBATED BY RISING SEA LEVELS AND INCREASED STORM INTENSITY, ALSO CONTRIBUTES TO HABITAT DEGRADATION.

SEAGRASS MEADOWS, ANOTHER CRITICAL HABITAT THAT PROVIDES FOOD AND SHELTER FOR NUMEROUS MARINE ANIMALS, ARE ALSO AFFECTED. INCREASED WATER DEPTH CAN REDUCE THE AMOUNT OF SUNLIGHT REACHING SEAGRASS, HINDERING PHOTOSYNTHESIS AND GROWTH. FURTHERMORE, CHANGES IN SALINITY DUE TO ALTERED FRESHWATER INFLOW AND INCREASED INUNDATION CAN STRESS SEAGRASS BEDS. THE LOSS OF THESE FOUNDATIONAL COASTAL HABITATS HAS PROFOUND IMPLICATIONS FOR MARINE BIODIVERSITY AND THE ECOLOGICAL SERVICES THEY PROVIDE, SUCH AS CARBON SEQUESTRATION AND SHORELINE PROTECTION.

CHANGES IN OCEAN CURRENTS AND NUTRIENT DISTRIBUTION

OCEAN CURRENTS ARE GLOBAL CONVEYOR BELTS THAT DISTRIBUTE HEAT, NUTRIENTS, AND OXYGEN THROUGHOUT THE WORLD'S OCEANS. CLIMATE CHANGE IS ALTERING THESE VITAL CIRCULATION PATTERNS, WITH SIGNIFICANT IMPLICATIONS FOR MARINE LIFE.

WARMING SURFACE WATERS CAN REDUCE THE MIXING BETWEEN SURFACE AND DEEP OCEAN LAYERS, A PROCESS CRUCIAL FOR BRINGING NUTRIENT-RICH WATERS FROM THE DEEP OCEAN TO THE SURFACE WHERE PHYTOPLANKTON CAN UTILIZE THEM. THIS REDUCTION IN NUTRIENT AVAILABILITY IN SURFACE WATERS CAN LEAD TO DECREASED PRIMARY PRODUCTIVITY, IMPACTING THE ENTIRE FOOD WEB. CONVERSELY, SOME REGIONS MAY EXPERIENCE INCREASED UPWELLING, BRINGING NUTRIENT-RICH, BUT POTENTIALLY WARMER OR MORE ACIDIC, WATERS TO THE SURFACE.

CHANGES IN OCEAN CURRENTS CAN ALSO ALTER THE DISTRIBUTION OF MARINE ORGANISMS. LARVAL STAGES OF MANY SPECIES DRIFT WITH CURRENTS, AND SHIFTS IN THESE PATTERNS CAN AFFECT WHERE THEY SETTLE AND THE CONNECTIVITY BETWEEN POPULATIONS. FURTHERMORE, ALTERED CURRENTS CAN INFLUENCE THE MIGRATION ROUTES OF LARGER MARINE ANIMALS, SUCH AS WHALES AND SEA TURTLES, IMPACTING THEIR ABILITY TO FIND FOOD AND BREEDING GROUNDS. THE COMPLEX INTERPLAY BETWEEN TEMPERATURE, SALINITY, AND CIRCULATION PATTERNS MAKES PREDICTING THE EXACT REGIONAL IMPACTS CHALLENGING BUT UNDENIABLY SIGNIFICANT.

IMPACTS ON MARINE BIODIVERSITY AND FOOD WEBS

THE CUMULATIVE EFFECTS OF RISING TEMPERATURES, ACIDIFICATION, SEA-LEVEL RISE, AND ALTERED CURRENTS ARE LEADING TO SIGNIFICANT SHIFTS IN MARINE BIODIVERSITY. SPECIES ARE FACING NEW ENVIRONMENTAL CHALLENGES, AND THEIR ABILITY TO ADAPT IS BEING TESTED. AS SOME SPECIES DECLINE OR SHIFT THEIR RANGES, OTHERS MAY THRIVE, LEADING TO NOVEL ECOLOGICAL COMMUNITIES AND ALTERED PREDATOR-PREY RELATIONSHIPS.

THE DELICATE BALANCE OF MARINE FOOD WEBS IS PARTICULARLY SUSCEPTIBLE TO DISRUPTION. FOR INSTANCE, THE DECLINE OF

KRILL POPULATIONS IN THE SOUTHERN OCEAN, LINKED TO WARMING WATERS AND REDUCED SEA ICE, HAS SIGNIFICANT IMPLICATIONS FOR THE WHALES, SEALS, AND PENGUINS THAT RELY ON THEM AS A PRIMARY FOOD SOURCE. SIMILARLY, THE LOSS OF CORAL REEFS REMOVES ESSENTIAL HABITAT AND FOOD SOURCES FOR THOUSANDS OF FISH SPECIES, LEADING TO A DECLINE IN REEF-ASSOCIATED BIODIVERSITY.

CHANGES IN THE TIMING OF BIOLOGICAL EVENTS, KNOWN AS PHENOLOGY, ARE ALSO OCCURRING. FOR EXAMPLE, THE SEASONAL BLOOMS OF PHYTOPLANKTON, THE BASE OF THE MARINE FOOD WEB, MAY OCCUR EARLIER OR LATER IN THE YEAR, CREATING A MISMATCH WITH THE LIFE CYCLES OF THE ZOOPLANKTON THAT FEED ON THEM, AND SUBSEQUENTLY AFFECTING HIGHER TROPHIC LEVELS. THESE CASCADING EFFECTS CAN LEAD TO SIGNIFICANT DECLINES IN POPULATION SIZES AND EVEN LOCAL EXTINCTIONS, FUNDAMENTALLY RESHAPING MARINE ECOSYSTEMS.

SPECIFIC VULNERABILITIES OF MARINE ECOSYSTEMS

WHILE ALL MARINE ECOSYSTEMS ARE AFFECTED BY CLIMATE CHANGE, SOME ARE INHERENTLY MORE VULNERABLE DUE TO THEIR SPECIFIC CHARACTERISTICS AND THE SPECIES THEY SUPPORT.

- **POLAR ECOSYSTEMS:** THE ARCTIC AND ANTARCTIC REGIONS ARE WARMING AT A FASTER RATE THAN THE GLOBAL AVERAGE. THIS LEADS TO RAPID MELTING OF SEA ICE, WHICH IS CRITICAL HABITAT FOR ICE-DEPENDENT SPECIES LIKE POLAR BEARS, SEALS, AND PENGUINS. THE LOSS OF SEA ICE ALSO IMPACTS THE PRODUCTIVITY OF THE UNDERLYING OCEAN, AFFECTING PHYTOPLANKTON AND ZOOPLANKTON POPULATIONS.
- **CORAL REEFS:** AS MENTIONED PREVIOUSLY, CORAL REEFS ARE EXTREMELY SENSITIVE TO TEMPERATURE CHANGES AND OCEAN ACIDIFICATION, MAKING THEM ONE OF THE MOST THREATENED ECOSYSTEMS ON EARTH. THEIR INTRICATE STRUCTURES PROVIDE HABITAT FOR A QUARTER OF ALL MARINE SPECIES.
- **ESTUARIES AND COASTAL WETLANDS:** THESE PRODUCTIVE ZONES ARE THREATENED BY SEA-LEVEL RISE, INCREASED STORM INTENSITY, AND CHANGES IN FRESHWATER INFLOW, IMPACTING THEIR SALINITY, BIODIVERSITY, AND ABILITY TO ACT AS NURSERIES FOR MARINE LIFE.
- **DEEP SEA ECOSYSTEMS:** WHILE SEEMINGLY REMOTE, DEEP-SEA ENVIRONMENTS ARE NOT IMMUNE. CHANGES IN OCEAN CIRCULATION CAN ALTER NUTRIENT DELIVERY, AND WARMING WATERS CAN IMPACT THE PHYSIOLOGY OF DEEP-SEA ORGANISMS ADAPTED TO STABLE, COLD CONDITIONS.

THE INTERCONNECTEDNESS OF THESE ECOSYSTEMS MEANS THAT THE DEGRADATION OF ONE CAN HAVE RIPPLE EFFECTS ON OTHERS, HIGHLIGHTING THE GLOBAL NATURE OF THE CLIMATE CHANGE THREAT TO MARINE LIFE.

CONSEQUENCES FOR HUMAN SOCIETIES

THE IMPACTS OF CLIMATE CHANGE ON MARINE LIFE HAVE PROFOUND CONSEQUENCES FOR HUMAN SOCIETIES WORLDWIDE. FISHERIES, A PRIMARY SOURCE OF PROTEIN AND LIVELIHOOD FOR BILLIONS OF PEOPLE, ARE DIRECTLY THREATENED BY SHIFTS IN FISH POPULATIONS, DECLINING CATCHES, AND DAMAGE TO FISHING INFRASTRUCTURE.

COASTAL COMMUNITIES FACE INCREASED RISKS FROM SEA-LEVEL RISE AND MORE FREQUENT AND INTENSE STORM SURGES, LEADING TO DISPLACEMENT, LOSS OF PROPERTY, AND ECONOMIC DISRUPTION. THE DEGRADATION OF CORAL REEFS AND OTHER COASTAL HABITATS REDUCES NATURAL DEFENSES AGAINST STORMS AND IMPACTS TOURISM INDUSTRIES THAT RELY ON HEALTHY MARINE ECOSYSTEMS.

FURTHERMORE, THE OCEAN PLAYS A CRUCIAL ROLE IN REGULATING GLOBAL CLIMATE AND SUPPORTING A VAST ARRAY OF BIODIVERSITY THAT HAS INTRINSIC VALUE AND PROVIDES POTENTIAL BENEFITS FOR MEDICINE AND BIOTECHNOLOGY. THE CONTINUED DEGRADATION OF MARINE ECOSYSTEMS DUE TO CLIMATE CHANGE JEOPARDIZES THESE ESSENTIAL SERVICES AND DIMINISHES THE PLANET'S NATURAL HERITAGE FOR FUTURE GENERATIONS.

FAQ

Q: HOW DOES RISING OCEAN TEMPERATURE DIRECTLY HARM MARINE ANIMALS?

A: RISING OCEAN TEMPERATURES CAN DISRUPT THE METABOLIC RATES, REPRODUCTIVE CYCLES, AND IMMUNE SYSTEMS OF MARINE ANIMALS. MANY SPECIES HAVE SPECIFIC TEMPERATURE RANGES THEY CAN TOLERATE, AND EXCEEDING THESE LIMITS CAN LEAD TO HEAT STRESS, REDUCED GROWTH, INCREASED SUSCEPTIBILITY TO DISEASES, AND EVEN DEATH. FOR EXAMPLE, FISH MAY STRUGGLE TO BREATHE IN WARMER, LESS OXYGENATED WATER, AND THE ENERGETIC DEMANDS OF MAINTAINING BODY TEMPERATURE CAN INCREASE SIGNIFICANTLY.

Q: WHAT IS OCEAN ACIDIFICATION AND WHY IS IT A PROBLEM FOR SHELLFISH?

A: OCEAN ACIDIFICATION IS THE PROCESS BY WHICH THE OCEAN'S pH DECREASES DUE TO THE ABSORPTION OF EXCESS ATMOSPHERIC CARBON DIOXIDE. THIS MAKES IT HARDER FOR MARINE ORGANISMS LIKE SHELLFISH, CORALS, AND CERTAIN PLANKTON TO BUILD AND MAINTAIN THEIR SHELLS AND SKELETONS, WHICH ARE MADE OF CALCIUM CARBONATE. AS THE OCEAN BECOMES LESS ALKALINE, THERE ARE FEWER CARBONATE IONS AVAILABLE, HINDERING CALCIFICATION AND POTENTIALLY CAUSING EXISTING SHELLS TO WEAKEN OR EVEN DISSOLVE.

Q: CAN SEA-LEVEL RISE AFFECT MARINE LIFE THAT LIVES FAR FROM THE COAST?

A: YES, SEA-LEVEL RISE CAN INDIRECTLY AFFECT MARINE LIFE FAR FROM THE COAST. IT LEADS TO INCREASED COASTAL EROSION, WHICH CAN RELEASE SEDIMENTS AND POLLUTANTS INTO THE OCEAN THAT TRAVEL AND IMPACT OFFSHORE ECOSYSTEMS. FURTHERMORE, THE LOSS OF COASTAL HABITATS LIKE MANGROVES AND SALT MARSHES, WHICH ACT AS IMPORTANT NURSERIES AND FEEDING GROUNDS FOR MANY SPECIES, REDUCES THE OVERALL HEALTH AND PRODUCTIVITY OF MARINE FOOD WEBS THAT EXTEND INTO DEEPER WATERS.

Q: HOW DO ALTERED OCEAN CURRENTS IMPACT THE DISTRIBUTION OF MARINE SPECIES?

A: OCEAN CURRENTS ACT AS HIGHWAYS FOR MARINE LIFE, TRANSPORTING LARVAE, NUTRIENTS, AND HEAT. CHANGES IN THESE CURRENTS CAN DISRUPT ESTABLISHED MIGRATION ROUTES, ALTER THE DISPERSAL PATTERNS OF PLANKTON AND JUVENILE FISH, AND CHANGE THE DELIVERY OF ESSENTIAL NUTRIENTS TO DIFFERENT REGIONS. THIS CAN LEAD TO SPECIES BEING FOUND IN NEW AREAS, STRUGGLING TO FIND FOOD OR SUITABLE BREEDING GROUNDS, OR EVEN BECOMING STRANDED IF CURRENTS SHIFT UNPREDICTABLY.

Q: WHAT IS CORAL BLEACHING AND WHAT CAUSES IT?

A: CORAL BLEACHING IS A PHENOMENON WHERE CORALS EXPEL THE SYMBIOTIC ALGAE (ZOOXANTHELLAE) THAT LIVE WITHIN THEIR TISSUES. THESE ALGAE PROVIDE CORALS WITH UP TO 90% OF THEIR ENERGY AND THEIR VIBRANT COLORS. BLEACHING IS PRIMARILY CAUSED BY INCREASES IN OCEAN TEMPERATURES, WHICH STRESS THE CORALS AND LEAD THEM TO EXPEL THEIR ALGAE. WHILE CORALS CAN RECOVER IF TEMPERATURES RETURN TO NORMAL QUICKLY, PROLONGED BLEACHING EVENTS CAN LEAD TO CORAL DEATH AND THE COLLAPSE OF REEF ECOSYSTEMS.

Q: ARE POLAR BEARS DIRECTLY THREATENED BY CLIMATE CHANGE EFFECTS ON MARINE LIFE?

A: YES, POLAR BEARS ARE DIRECTLY THREATENED BY CLIMATE CHANGE EFFECTS ON MARINE LIFE, PARTICULARLY THROUGH THE LOSS OF ARCTIC SEA ICE. SEA ICE IS THEIR PRIMARY PLATFORM FOR HUNTING SEALS, THEIR MAIN FOOD SOURCE. AS THE ICE MELTS EARLIER IN THE SPRING AND FORMS LATER IN THE FALL, POLAR BEARS HAVE LESS TIME TO HUNT, LEADING TO THINNER BEARS, REDUCED REPRODUCTIVE SUCCESS, AND INCREASED MORTALITY. THE CHANGES IN SEA ICE ALSO AFFECT THE DISTRIBUTION AND AVAILABILITY OF SEALS.

Q: HOW DOES OCEAN ACIDIFICATION IMPACT THE ENTIRE MARINE FOOD WEB?

A: OCEAN ACIDIFICATION HAS A RIPPLE EFFECT THROUGHOUT THE MARINE FOOD WEB. ORGANISMS AT THE BASE OF THE FOOD WEB, SUCH AS CALCIFYING PLANKTON (LIKE PTEROPODS), ARE DIRECTLY AFFECTED BY THE DIFFICULTY IN FORMING SHELLS. A DECLINE IN THESE FOUNDATIONAL SPECIES CAN LEAD TO FOOD SHORTAGES FOR THE ZOOPLANKTON THAT FEED ON THEM, AND SUBSEQUENTLY FOR THE FISH, SEABIRDS, AND MARINE MAMMALS THAT RELY ON THOSE ZOOPLANKTON. THIS DISRUPTION AT THE BASE CAN WEAKEN THE ENTIRE ECOSYSTEM.

Q: WHAT ARE THE ECONOMIC CONSEQUENCES OF CLIMATE CHANGE IMPACTS ON MARINE LIFE?

A: THE ECONOMIC CONSEQUENCES ARE SIGNIFICANT AND WIDESPREAD. THEY INCLUDE LOSSES IN FISHERIES REVENUE DUE TO DECLINING FISH STOCKS AND SHIFTS IN SPECIES DISTRIBUTION, REDUCED INCOME FROM TOURISM AS CORAL REEFS AND OTHER MARINE ATTRACTIONS DEGRADE, AND INCREASED COSTS ASSOCIATED WITH COASTAL PROTECTION AND DISASTER RECOVERY FROM INTENSIFIED STORMS. FURTHERMORE, THE LOSS OF ECOSYSTEM SERVICES, SUCH AS CARBON SEQUESTRATION BY MARINE ORGANISMS, HAS UNQUANTIFIED BUT SUBSTANTIAL ECONOMIC IMPLICATIONS.

Climate Change Effects Marine Life

Climate Change Effects Marine Life

Related Articles

- [clinical anatomy pt](#)
- [claudé monet patronage](#)
- [climate change impact on resilient cities](#)

[Back to Home](#)