

DARK MATTER CHARACTERISTICS

UNVEILING THE MYSTERIES: A DEEP DIVE INTO DARK MATTER CHARACTERISTICS

DARK MATTER CHARACTERISTICS ARE AMONG THE MOST PROFOUND ENIGMAS CONFRONTING MODERN COSMOLOGY. WHILE INVISIBLE TO OUR TELESCOPES AND UNDETECTABLE THROUGH ELECTROMAGNETIC RADIATION, ITS GRAVITATIONAL INFLUENCE IS UNDENIABLY SHAPING THE VERY FABRIC OF THE UNIVERSE. FROM THE STATELY DANCE OF GALAXIES TO THE GRAND COSMIC WEB, DARK MATTER PLAYS A PIVOTAL ROLE, YET ITS FUNDAMENTAL NATURE REMAINS ELUSIVE. THIS ARTICLE WILL EMBARK ON A COMPREHENSIVE EXPLORATION OF WHAT WE CURRENTLY UNDERSTAND ABOUT ITS PROPERTIES, HOW SCIENTISTS INFER ITS EXISTENCE, AND THE ONGOING QUEST TO DIRECTLY DETECT THIS MYSTERIOUS SUBSTANCE. WE WILL DELVE INTO ITS GRAVITATIONAL EFFECTS, ITS LACK OF INTERACTION WITH LIGHT, AND THE LEADING THEORETICAL CANDIDATES THAT MIGHT EXPLAIN ITS EXISTENCE, ALL WHILE PAINTING A DETAILED PICTURE OF THIS UNSEEN COSMIC ARCHITECT.

TABLE OF CONTENTS

WHAT IS DARK MATTER?

KEY CHARACTERISTICS OF DARK MATTER

INFERRING DARK MATTER'S PRESENCE

LEADING CANDIDATES FOR DARK MATTER

THE SEARCH FOR DIRECT DETECTION

WHAT IS DARK MATTER?

UNDERSTANDING DARK MATTER BEGINS WITH ACKNOWLEDGING ITS UNIQUE STATUS IN THE COSMIC INVENTORY. IT IS NOT "DARK" IN THE SENSE OF BEING MERELY UNLIT MATTER LIKE A COLD PLANET; RATHER, IT IS FUNDAMENTALLY NON-LUMINOUS, MEANING IT DOES NOT EMIT, ABSORB, OR REFLECT ANY FORM OF ELECTROMAGNETIC RADIATION. THIS INCLUDES VISIBLE LIGHT, RADIO WAVES, X-RAYS, AND GAMMA RAYS. THIS FUNDAMENTAL CHARACTERISTIC IS WHAT RENDERS IT INVISIBLE TO CONVENTIONAL ASTRONOMICAL INSTRUMENTS. DESPITE ITS INVISIBILITY, ITS GRAVITATIONAL PULL IS A DOMINANT FORCE, ACCOUNTING FOR APPROXIMATELY 85% OF THE TOTAL MATTER IN THE UNIVERSE. THE REMAINING 15% IS WHAT WE PERCEIVE AS "NORMAL" OR BARYONIC MATTER, THE STUFF STARS, PLANETS, AND WE ARE MADE OF. THE CONCEPT OF DARK MATTER EMERGED FROM DISCREPANCIES OBSERVED IN GALACTIC ROTATION CURVES AND THE DYNAMICS OF GALAXY CLUSTERS, WHERE VISIBLE MATTER ALONE COULD NOT EXPLAIN THE OBSERVED GRAVITATIONAL EFFECTS.

KEY CHARACTERISTICS OF DARK MATTER

THE DEFINING ATTRIBUTES OF DARK MATTER ARE DERIVED FROM ITS OBSERVED GRAVITATIONAL INTERACTIONS AND ITS LACK OF ENGAGEMENT WITH THE ELECTROMAGNETIC SPECTRUM. THESE CHARACTERISTICS ARE CRUCIAL FOR DISTINGUISHING IT FROM ORDINARY MATTER AND FOR GUIDING THEORETICAL MODELS.

GRAVITATIONAL INFLUENCE

PERHAPS THE MOST COMPELLING CHARACTERISTIC OF DARK MATTER IS ITS POWERFUL GRAVITATIONAL PULL. WE INFER ITS EXISTENCE PRECISELY BECAUSE OF THE GRAVITATIONAL FORCES IT EXERTS ON VISIBLE MATTER. WITHOUT DARK MATTER, GALAXIES WOULD SPIN THEMSELVES APART, AND GALAXY CLUSTERS WOULD NOT HOLD TOGETHER. THINK OF IT LIKE AN INVISIBLE SCAFFOLDING THAT PROVIDES THE NECESSARY GRAVITATIONAL GLUE FOR COSMIC STRUCTURES TO FORM AND PERSIST. THIS GRAVITATIONAL DOMINANCE IS EVIDENT ON SCALES RANGING FROM INDIVIDUAL GALAXIES TO THE LARGEST OBSERVABLE STRUCTURES IN THE UNIVERSE.

Non-Baryonic Nature

A CRITICAL CHARACTERISTIC IS THAT DARK MATTER IS OVERWHELMINGLY NON-BARYONIC. BARYONIC MATTER IS COMPOSED OF PROTONS AND NEUTRONS, THE BUILDING BLOCKS OF ATOMS. OBSERVATIONS OF BIG BANG NUCLEOSYNTHESIS, WHICH PRECISELY PREDICT THE ABUNDANCES OF LIGHT ELEMENTS LIKE HYDROGEN AND HELIUM FORMED IN THE EARLY UNIVERSE, LIMIT THE TOTAL AMOUNT OF BARYONIC MATTER TO ONLY ABOUT 5% OF THE UNIVERSE'S TOTAL ENERGY DENSITY. SINCE WE OBSERVE GRAVITATIONAL EFFECTS EQUIVALENT TO ABOUT 27% OF THE UNIVERSE'S ENERGY DENSITY IN THE FORM OF MATTER, THE VAST MAJORITY OF THIS MATTER MUST BE SOMETHING ELSE ENTIRELY – HENCE, NON-BARYONIC DARK MATTER.

Weak Interaction with Itself and Ordinary Matter

ANOTHER CRUCIAL ASPECT OF DARK MATTER IS ITS EXTREMELY WEAK INTERACTION WITH ITSELF AND WITH ORDINARY BARYONIC MATTER, APART FROM GRAVITY. THIS MEANS IT DOESN'T READILY FORM CLOUDS, COALESCE INTO STARS, OR INTERACT THROUGH ELECTROMAGNETIC FORCES. IF DARK MATTER PARTICLES COULD COLLIDE AND INTERACT ELECTROMAGNETICALLY, WE WOULD EXPECT TO SEE EVIDENCE OF THIS IN THE FORM OF RADIATION OR OTHER OBSERVABLE PHENOMENA, WHICH WE DO NOT. THIS "COLLISIONLESS" NATURE IS A KEY ASSUMPTION IN MANY COSMOLOGICAL SIMULATIONS.

Cold or Warm Nature

THE TEMPERATURE AND VELOCITY OF DARK MATTER PARTICLES IN THE EARLY UNIVERSE ARE SIGNIFICANT. COSMOLOGISTS GENERALLY CATEGORIZE DARK MATTER AS EITHER "COLD" OR "WARM." COLD DARK MATTER (CDM) PARTICLES WERE MOVING RELATIVELY SLOWLY IN THE EARLY UNIVERSE, ALLOWING FOR THE HIERARCHICAL FORMATION OF STRUCTURE, WHERE SMALL STRUCTURES MERGE TO FORM LARGER ONES. THIS IS CONSISTENT WITH OBSERVATIONS OF THE COSMIC WEB AND GALAXY FORMATION. WARM DARK MATTER (WDM) PARTICLES WOULD HAVE MOVED FASTER, POTENTIALLY SUPPRESSING THE FORMATION OF THE SMALLEST STRUCTURES. CURRENT COSMOLOGICAL MODELS STRONGLY FAVOR CDM, THOUGH SOME WDM SCENARIOS ARE STILL BEING EXPLORED.

Stability

IT IS ASSUMED THAT DARK MATTER PARTICLES ARE STABLE, OR AT LEAST POSSESS AN EXTREMELY LONG HALF-LIFE, MEANING THEY DO NOT DECAY INTO LIGHTER PARTICLES OVER COSMOLOGICAL TIMESCALES. IF DARK MATTER WERE TO DECAY RAPIDLY, ITS GRAVITATIONAL INFLUENCE WOULD HAVE DIMINISHED OVER COSMIC HISTORY, WHICH IS NOT WHAT WE OBSERVE. ITS PERSISTENT GRAVITATIONAL PRESENCE SUGGESTS IT IS A FUNDAMENTAL AND ENDURING COMPONENT OF THE UNIVERSE.

Inferring Dark Matter's Presence

SINCE WE CANNOT SEE DARK MATTER DIRECTLY, ITS EXISTENCE IS INFERRED THROUGH A VARIETY OF INDIRECT ASTRONOMICAL OBSERVATIONS THAT REVEAL ITS GRAVITATIONAL EFFECTS. THESE METHODS PROVIDE COMPELLING EVIDENCE FOR ITS PRESENCE AND ALLOW US TO MAP ITS DISTRIBUTION.

Galactic Rotation Curves

ONE OF THE EARLIEST AND MOST ROBUST PIECES OF EVIDENCE FOR DARK MATTER COMES FROM STUDYING HOW GALAXIES ROTATE. STARS AND GAS CLOUDS ORBITING THE OUTER EDGES OF SPIRAL GALAXIES MOVE MUCH FASTER THAN PREDICTED BASED ON THE VISIBLE MASS ALONE. ACCORDING TO NEWTONIAN GRAVITY, ORBITAL SPEED SHOULD DECREASE WITH DISTANCE FROM THE GALACTIC CENTER, SIMILAR TO HOW PLANETS FARTHER FROM THE SUN ORBIT MORE SLOWLY. HOWEVER, OBSERVATIONS SHOW THAT THE ORBITAL SPEEDS REMAIN REMARKABLY CONSTANT OR EVEN INCREASE SLIGHTLY AT LARGER RADII. THIS "FLAT" ROTATION CURVE STRONGLY SUGGESTS THE PRESENCE OF AN UNSEEN HALO OF MATTER EXTENDING FAR BEYOND THE VISIBLE DISK

OF THE GALAXY, PROVIDING THE EXTRA GRAVITATIONAL PULL.

GRAVITATIONAL LENSING

GRAVITATIONAL LENSING IS ANOTHER POWERFUL TOOL FOR DETECTING AND MAPPING DARK MATTER. MASSIVE OBJECTS, INCLUDING CONCENTRATIONS OF DARK MATTER, WARP THE FABRIC OF SPACETIME. LIGHT FROM DISTANT OBJECTS PASSING THROUGH THIS WARPED SPACETIME IS BENT, MUCH LIKE LIGHT PASSING THROUGH A LENS. THIS BENDING CAN DISTORT THE IMAGES OF BACKGROUND GALAXIES, MAGNIFY THEM, OR EVEN CREATE MULTIPLE IMAGES OF THE SAME OBJECT. BY ANALYZING THE DEGREE AND PATTERN OF THIS DISTORTION, ASTRONOMERS CAN MAP THE DISTRIBUTION OF MASS, INCLUDING DARK MATTER, IN FOREGROUND GALAXY CLUSTERS AND OTHER LARGE-SCALE STRUCTURES. THE BULLET CLUSTER, FOR INSTANCE, PROVIDES A STUNNING VISUAL DEMONSTRATION OF DARK MATTER'S SEPARATION FROM NORMAL MATTER DURING A COSMIC COLLISION.

COSMIC MICROWAVE BACKGROUND (CMB) RADIATION

THE FAINT AFTERGLOW OF THE BIG BANG, THE COSMIC MICROWAVE BACKGROUND RADIATION, ALSO HOLDS CRUCIAL INFORMATION ABOUT DARK MATTER. TINY TEMPERATURE FLUCTUATIONS IN THE CMB REVEAL THE DENSITY VARIATIONS IN THE EARLY UNIVERSE. THE PATTERN AND AMPLITUDE OF THESE FLUCTUATIONS ARE SENSITIVE TO THE RELATIVE AMOUNTS OF BARYONIC MATTER, DARK MATTER, AND DARK ENERGY. PRECISE MEASUREMENTS OF THE CMB BY MISSIONS LIKE WMAP AND PLANCK HAVE ALLOWED COSMOLOGISTS TO CONSTRAIN THE COMPOSITION OF THE UNIVERSE WITH REMARKABLE ACCURACY, CONFIRMING THE DOMINANCE OF DARK MATTER.

LARGE-SCALE STRUCTURE FORMATION

THE DISTRIBUTION OF GALAXIES AND GALAXY CLUSTERS THROUGHOUT THE UNIVERSE, KNOWN AS THE LARGE-SCALE STRUCTURE, IS ANOTHER KEY PIECE OF EVIDENCE. COMPUTER SIMULATIONS THAT AIM TO REPRODUCE THE OBSERVED COSMIC WEB, INCLUDING THE FILAMENTARY STRUCTURES AND VOIDS, REQUIRE THE PRESENCE OF DARK MATTER TO SEED THE INITIAL DENSITY FLUCTUATIONS THAT GRAVITY THEN AMPLIFIES. WITHOUT DARK MATTER'S GRAVITATIONAL INFLUENCE, THE UNIVERSE WOULD LIKELY BE MUCH MORE UNIFORM AND LACK THE COMPLEX STRUCTURES WE OBSERVE TODAY.

LEADING CANDIDATES FOR DARK MATTER

THE ELUSIVE NATURE OF DARK MATTER HAS LED TO A PLETHORA OF THEORETICAL PROPOSALS FOR WHAT IT MIGHT BE. SCIENTISTS ARE EXPLORING VARIOUS TYPES OF PARTICLES, EACH WITH ITS OWN SET OF PREDICTED PROPERTIES.

WIMPS (WEAKLY INTERACTING MASSIVE PARTICLES)

FOR A LONG TIME, WIMPS HAVE BEEN A LEADING CANDIDATE FOR DARK MATTER. THESE HYPOTHETICAL PARTICLES ARE THEORIZED TO HAVE MASSES ROUGHLY BETWEEN A FEW GeV AND A FEW TeV (GIGAELECTRONVOLTS TO TERA-ELECTRONVOLTS) AND TO INTERACT ONLY WEAKLY WITH ORDINARY MATTER AND THEMSELVES, PRIMARILY THROUGH GRAVITY AND THE WEAK NUCLEAR FORCE. THIS WEAK INTERACTION ALIGNS WITH THE OBSERVED PROPERTIES OF DARK MATTER. MANY DIRECT DETECTION EXPERIMENTS ARE SPECIFICALLY DESIGNED TO HUNT FOR WIMPS BY LOOKING FOR THE RARE INSTANCES WHERE A WIMP MIGHT COLLIDE WITH AN ATOMIC NUCLEUS IN A DETECTOR.

AXIONS

AXIONS ARE ANOTHER PROMISING CANDIDATE, ARISING FROM SOLUTIONS TO A PROBLEM IN QUANTUM CHROMODYNAMICS

(QCD). They are predicted to be extremely light particles, much lighter than WIMPs, and to interact very weakly with other matter. Their weak interaction makes them incredibly difficult to detect directly, but their potential abundance makes them a significant contender for making up a portion or even all of dark matter. Experiments are underway to search for axions by looking for their potential conversion into photons in strong magnetic fields.

STERILE NEUTRINOS

Neutrinos are known particles that interact only via the weak force and gravity. Standard neutrinos are too light and fast-moving to account for the observed dark matter. However, theoretical extensions to the Standard Model propose the existence of "sterile neutrinos," which would interact even more weakly than standard neutrinos and could potentially have the right mass and abundance to be dark matter. Detecting sterile neutrinos would be an extraordinary feat due to their extremely feeble interactions.

PRIMORDIAL BLACK HOLES

While less favored than particle candidates, primordial black holes (PBHs) are another possibility. These are hypothetical black holes that could have formed in the extremely dense conditions of the early universe, shortly after the Big Bang. If they exist in the right mass range and abundance, they could contribute to or entirely comprise dark matter. However, observational constraints from gravitational lensing and other astrophysical phenomena have significantly narrowed the possible mass windows for PBHs to constitute all of dark matter.

THE SEARCH FOR DIRECT DETECTION

The quest to directly detect dark matter particles is one of the most active frontiers in physics and astronomy. Numerous experiments are underway, employing diverse strategies to capture these elusive particles.

UNDERGROUND DETECTORS

Many experiments involve highly sensitive detectors placed deep underground. This is to shield them from cosmic rays and other sources of background radiation that could mimic a dark matter signal. These detectors often use large volumes of super-pure materials, such as noble liquids (like Xenon or Argon) or crystalline solids, designed to register the faint recoil energy imparted when a dark matter particle collides with an atomic nucleus. Examples include experiments like LUX-ZEPLIN (LZ), XENONnT, and SuperCDMS.

PARTICLE ACCELERATORS

Particle accelerators, such as the Large Hadron Collider (LHC) at CERN, are also involved in the search for dark matter. By smashing protons together at incredibly high energies, scientists hope to produce dark matter particles that would then escape the detector unseen. Their presence would be inferred from missing energy and momentum in the collision debris, suggesting that some invisible particles carried that energy away. This indirect approach complements direct detection efforts by searching for potential dark matter constituents created in laboratory conditions.

SATELLITE AND SPACE-BASED EXPERIMENTS

SOME EXPERIMENTS LOOK FOR INDIRECT SIGNS OF DARK MATTER ANNIHILATION OR DECAY. IF DARK MATTER PARTICLES CAN COLLIDE AND ANNIHILATE EACH OTHER, THEY COULD PRODUCE DETECTABLE PARTICLES LIKE GAMMA RAYS, POSITRONS, OR NEUTRINOS. SPACE TELESCOPES LIKE THE FERMI GAMMA-RAY SPACE TELESCOPE AND EXPERIMENTS ON SATELLITES ARE DESIGNED TO SEARCH FOR AN EXCESS OF THESE PARTICLES COMING FROM REGIONS WHERE DARK MATTER IS EXPECTED TO BE DENSE, SUCH AS THE GALACTIC CENTER OR DWARF GALAXIES.

THE ONGOING INVESTIGATION INTO DARK MATTER CHARACTERISTICS CONTINUES TO PUSH THE BOUNDARIES OF OUR UNDERSTANDING OF THE COSMOS. EACH NEW OBSERVATION AND EXPERIMENTAL RESULT BRINGS US CLOSER TO UNRAVELING THIS PROFOUND MYSTERY, POTENTIALLY LEADING TO NEW PHYSICS BEYOND OUR CURRENT STANDARD MODEL AND A MORE COMPLETE PICTURE OF OUR UNIVERSE'S COMPOSITION AND EVOLUTION.

Q: WHAT IS THE PRIMARY EVIDENCE FOR THE EXISTENCE OF DARK MATTER?

A: THE PRIMARY EVIDENCE FOR DARK MATTER'S EXISTENCE COMES FROM ITS OBSERVED GRAVITATIONAL EFFECTS ON VISIBLE MATTER. THIS INCLUDES DISCREPANCIES IN GALACTIC ROTATION CURVES, WHERE STARS ORBIT FASTER THAN EXPECTED BASED ON VISIBLE MASS ALONE, AND THE GRAVITATIONAL LENSING OF LIGHT BY MASSIVE STRUCTURES LIKE GALAXY CLUSTERS, WHICH REVEALS MORE MASS THAN CAN BE ACCOUNTED FOR BY NORMAL MATTER. ADDITIONALLY, THE PATTERNS IN THE COSMIC MICROWAVE BACKGROUND RADIATION AND THE FORMATION OF LARGE-SCALE COSMIC STRUCTURES ARE BEST EXPLAINED BY THE PRESENCE OF A SIGNIFICANT AMOUNT OF UNSEEN MATTER.

Q: HOW DOES DARK MATTER DIFFER FROM NORMAL BARYONIC MATTER?

A: DARK MATTER DIFFERS FUNDAMENTALLY FROM NORMAL BARYONIC MATTER IN THAT IT DOES NOT INTERACT WITH ELECTROMAGNETIC RADIATION. THIS MEANS IT DOES NOT EMIT, ABSORB, OR REFLECT LIGHT, MAKING IT INVISIBLE TO TELESCOPES. WHILE BARYONIC MATTER IS COMPOSED OF PROTONS AND NEUTRONS, FORMING ATOMS AND MOLECULES, DARK MATTER IS BELIEVED TO BE COMPOSED OF EXOTIC PARTICLES THAT INTERACT PRIMARILY THROUGH GRAVITY AND POSSIBLY THE WEAK NUCLEAR FORCE.

Q: WHY IS DARK MATTER CALLED "DARK"?

A: DARK MATTER IS CALLED "DARK" BECAUSE IT IS INVISIBLE TO US. IT DOES NOT EMIT, ABSORB, OR REFLECT ANY FORM OF LIGHT OR OTHER ELECTROMAGNETIC RADIATION, WHICH ARE THE PRIMARY WAYS ASTRONOMERS DETECT AND STUDY CELESTIAL OBJECTS. ITS PRESENCE IS ONLY INFERRED THROUGH ITS GRAVITATIONAL INFLUENCE.

Q: WHAT ARE THE MAIN PROPERTIES SCIENTISTS ATTRIBUTE TO DARK MATTER BASED ON CURRENT OBSERVATIONS?

A: BASED ON CURRENT OBSERVATIONS, SCIENTISTS ATTRIBUTE SEVERAL KEY PROPERTIES TO DARK MATTER. THESE INCLUDE ITS SIGNIFICANT GRAVITATIONAL INFLUENCE, ITS NON-BARYONIC NATURE (MEANING IT'S NOT MADE OF PROTONS AND NEUTRONS), ITS WEAK INTERACTION WITH ITSELF AND ORDINARY MATTER (APART FROM GRAVITY), ITS PRESUMED STABILITY OVER COSMIC TIMESCALES, AND ITS LIKELY "COLD" NATURE, MEANING IT WAS MOVING SLOWLY IN THE EARLY UNIVERSE, FACILITATING STRUCTURE FORMATION.

Q: WHAT ARE WIMPS AND WHY ARE THEY CONSIDERED A CANDIDATE FOR DARK MATTER?

A: WIMPS, WHICH STANDS FOR WEAKLY INTERACTING MASSIVE PARTICLES, ARE HYPOTHETICAL PARTICLES THAT ARE CONSIDERED A LEADING CANDIDATE FOR DARK MATTER. THEY ARE THEORIZED TO HAVE MASSES SIGNIFICANTLY GREATER THAN

PROTONS AND TO INTERACT WITH ORDINARY MATTER AND THEMSELVES ONLY VERY WEAKLY, PRIMARILY THROUGH GRAVITY AND THE WEAK NUCLEAR FORCE. THIS WEAK INTERACTION ALIGNS WITH THE OBSERVED BEHAVIOR OF DARK MATTER, WHICH EXERTS GRAVITATIONAL INFLUENCE WITHOUT PARTICIPATING IN ELECTROMAGNETIC INTERACTIONS.

Q: HOW ARE SCIENTISTS ATTEMPTING TO DIRECTLY DETECT DARK MATTER PARTICLES?

A: SCIENTISTS ARE ATTEMPTING TO DIRECTLY DETECT DARK MATTER PARTICLES THROUGH VARIOUS EXPERIMENTAL METHODS. MANY EXPERIMENTS USE HIGHLY SENSITIVE DETECTORS, OFTEN PLACED DEEP UNDERGROUND TO SHIELD THEM FROM BACKGROUND RADIATION, DESIGNED TO REGISTER THE FAINT RECOIL ENERGY WHEN A HYPOTHETICAL DARK MATTER PARTICLE COLLIDES WITH AN ATOMIC NUCLEUS. OTHER APPROACHES INVOLVE SEARCHING FOR INDIRECT EVIDENCE, SUCH AS THE PRODUCTS OF DARK MATTER ANNIHILATION OR DECAY, USING PARTICLE ACCELERATORS OR SPACE-BASED TELESCOPES.

Q: WHAT IS GRAVITATIONAL LENSING AND HOW DOES IT PROVIDE EVIDENCE FOR DARK MATTER?

A: GRAVITATIONAL LENSING IS A PHENOMENON WHERE THE GRAVITY OF A MASSIVE OBJECT, SUCH AS A GALAXY OR GALAXY CLUSTER, BENDS THE PATH OF LIGHT FROM MORE DISTANT OBJECTS. THIS BENDING CAN DISTORT, MAGNIFY, OR CREATE MULTIPLE IMAGES OF THE BACKGROUND OBJECTS. BY ANALYZING THE DEGREE AND PATTERN OF THIS LENSING EFFECT, ASTRONOMERS CAN MAP THE DISTRIBUTION OF MASS, INCLUDING DARK MATTER, IN THE FOREGROUND STRUCTURE. THE OBSERVED LENSING EFFECTS ARE OFTEN FAR STRONGER THAN CAN BE EXPLAINED BY THE VISIBLE MATTER ALONE, THUS PROVIDING COMPELLING EVIDENCE FOR THE PRESENCE OF DARK MATTER.

Q: CAN DARK MATTER BE MADE OF BLACK HOLES?

A: THE POSSIBILITY THAT DARK MATTER COULD BE COMPOSED OF PRIMORDIAL BLACK HOLES (PBHS) HAS BEEN CONSIDERED. THESE ARE HYPOTHETICAL BLACK HOLES THAT COULD HAVE FORMED IN THE VERY EARLY UNIVERSE. WHILE PBHS COULD THEORETICALLY CONTRIBUTE TO DARK MATTER, OBSERVATIONAL CONSTRAINTS FROM GRAVITATIONAL LENSING AND OTHER ASTROPHYSICAL PHENOMENA HAVE SIGNIFICANTLY LIMITED THE MASS RANGES AND ABUNDANCES FOR WHICH THEY COULD CONSTITUTE ALL OF THE UNIVERSE'S DARK MATTER. PARTICLE CANDIDATES ARE CURRENTLY CONSIDERED MORE LIKELY.

Dark Matter Characteristics

Dark Matter Characteristics

Related Articles

- [cybersecurity policy united states](#)
- [cyberbullying prevention strategies for legal professionals](#)
- [d-day impact on us economy](#)

[Back to Home](#)